

Vortex Tiling in a Spin-2 Spinor Bose-Einstein Condensate

Michikazu Kobayashi¹, Yuki Kawaguchi¹, Masahito Ueda^{1,2}

¹*Department of Physics, University of Tokyo, Hongo 7-3-1, Bunkyo-ku, Tokyo 113-0033, Japan. and*

²*ERATO Macroscopic Quantum Control Project, JST, Tokyo 113-8656, Japan*

(Dated: July 21, 2009)

We investigate core structures of fractional vortices in a spin-2 spinor Bose-Einstein condensate and find that commensurability between discrete symmetries of the order parameters inside and outside of a vortex core brings about spontaneous breaking of the rotational symmetry of the core and the associated twisting excitations. Our predictions can be tested experimentally by using a rotating spin-2 ⁸⁷Rb Bose-Einstein condensate and enable an experimental realization of the cyclic order parameter, regardless of whether the ground state is cyclic or otherwise.

PACS numbers: 05.30.Jp, 03.75.Lm, 03.75.Mn, 11.27.+d

Quantized vortices are topological defects of the superfluid order parameter. For single component Bose-Einstein condensates (BECs), the quantized vortices emerge as phase singularities, and the order parameter vanishes at the core center. However, the situation drastically changes for spinor BECs, where the vortex core can be filled with a state having different symmetry. For example, a spin-1 polar BEC supports a half-quantized vortex with a ferromagnetic core [1], and a spin-1 ferromagnetic BEC can hold a spin vortex with a polar core. The vortex core structure for the spin-1 BEC is, however, uniquely determined since there are only two possible ground-state phases in zero magnetic field. The problem becomes much more involved in higher-spin BECs because of the greater possibilities of the ground-state phases. Since the order parameter of the spinor BEC is generally anisotropic, the following questions arise which we refer to as “vortex tiling problems”: (1) which state fills the vortex core?; (2) which orientation the order parameter of the vortex core takes with respect to the surrounding BEC?; (3) how can the vortex core be embedded in the BEC if they have different discrete symmetries? We will show that the commensurability in spin space gives rise to breaking of the rotational symmetry of the vortex in real space, thus allowing us to probe the symmetry of the order parameter by observing the deformed shape of the vortex core.

In this Letter, we address the tiling problems for fractional vortices in a spin-2 BEC. The ground state of the spin-2 BEC is shown to have four different phases under zero magnetic field [2, 3, 4, 5, 6]; ferromagnetic, uniaxial nematic, biaxial nematic, and cyclic, where the last three phases have discrete symmetries. The biaxial nematic phase has the non-Abelian fourth dihedral (D_4) symmetry [6], and the cyclic phase has the non-Abelian tetrahedral symmetry [7]. We consider here two highly nontrivial cases; a vortex in the biaxial nematic phase with the cyclic core, and a vortex in the cyclic phase with the biaxial nematic core. The solutions of these tiling problems are shown in Fig. 1. Figures 1 (a) and (b) show the cloverleaf and triad profiles of the biaxial




FIG. 1: (Color) (a) and (b) Plots of $|\Phi(\theta, \phi)|^2$ for the biaxial nematic phase Ψ_B and the cyclic phase Ψ_C , where $\Phi(\theta, \phi) \equiv \sum_{m=-2}^2 Y_{2m}(\theta, \phi) \psi_m$. Colors represent the phase of $\Phi(\theta, \phi)$. (c) and (d) Distributions of $\Phi(\theta, \phi)$ for the $1/2-1/4$ vortex with the cyclic core and the $1/3-1/3$ vortex with the biaxial nematic core. Here $1/2$ and $1/4$ refer to the π gauge transformation and the $\pi/2$ spin rotation around the vortex, respectively. (e) and (f) Profiles of $|A_{30}|^2$ and $|A_{20}|^2$ for the $1/2-1/4$ and $1/3-1/3$ vortices, where the size of each panel is 16×16 in units of the healing length $\xi = \hbar/\sqrt{2Mc_2 n_{\text{tot}}}$.

ial nematic and cyclic order parameters in spin space. The direction of the triad (cloverleaf) inside the vortex must be determined so as to be smoothly connected with the cloverleaf (triad) of the BECs outside of the vortex; Fig. 1 (c) (Fig. 1 (d)) shows the tiling pattern of the order parameters for these vortices. Each lobe of the triad (cloverleaf) in the core is connected with that of the cloverleaf (triad) outside the core. The smoothly connected directions are indicated by arrows with colors of the corresponding lobes. As a consequence, the rotational symmetry of the vortex core is spontaneously broken and the core is deformed to become triangular (elliptic) for the vortex in the biaxial nematic (cyclic) phase with the cyclic (biaxial nematic) core, as shown in Fig. 1 (e) (Fig. 1 (f)). Here the three-fold and two-fold

symmetries of the cores reflect those of the cyclic and biaxial nematic phases, respectively. Details of Fig. 1 are explained below.

Another important finding of this Letter is that the cyclic state can be realized as the vortex core state in an $F = 2$ ^{87}Rb BEC [8, 9, 10, 11]. Although the ground state of this BEC has yet to be determined between the cyclic and biaxial nematic phases due to the complications arising from the quadratic Zeeman effect and hyperfine-spin-exchanging relaxations, we have confirmed that the vortex with the cyclic core can be generated by rotating the biaxial nematic initial configuration of the $F = 2$ ^{87}Rb BEC, regardless of which phase the true ground state is. The core deformation predicted in this Letter can also be used as a hallmark of the cyclic state.

We consider a BEC of spin-2 atoms with mass M whose mean-field energy functional is given by [2, 3]

$$H = \int d\mathbf{x} \left[\sum_{m=-2}^2 \psi_m^* \left(-\frac{\hbar^2}{2M} \nabla^2 \right) \psi_m + \frac{c_0}{2} n_{\text{tot}}^2 + \frac{c_1}{2} |\mathbf{F}|^2 + \frac{c_2}{2} |A_{20}|^2 \right], \quad (1)$$

where ψ_m is the order parameter of the BEC in a magnetic sublevel $m = 0, \pm 1, \pm 2$ at position $\mathbf{r}$, and $n_{\text{tot}} = \sum_{m=-2}^2 |\psi_m|^2$, $\mathbf{F} = \sum_{m,m'=-2}^2 \psi_m^* \mathbf{f}_{m,m'} \psi_{m'}$, and $A_{20} = (2\psi_2\psi_{-2} - 2\psi_1\psi_{-1} + \psi_0^2)/\sqrt{5}$ are the total number density, the spin density, and the spin-singlet pair amplitude, respectively, with $\mathbf{f}_{m,m'}$ being a vector of spin-2 matrices. The ground state of Hamiltonian (1) is (i) ferromagnetic ($\mathbf{F} \neq 0$ and $A_{20} = 0$) for $c_1 < 0$ and $c_2 > 20c_1$, (ii) uniaxial nematic or biaxial nematic ($\mathbf{F} = 0$ and $A_{20} \neq 0$) for $c_2 < 0$ and $c_2 < 20c_1$, and (iii) cyclic ($\mathbf{F} = A_{20} = 0$) for $c_1 > 0$ and $c_2 > 0$. A representative order parameter for each phase is given by (i) $\Psi_F = \sqrt{n_{\text{tot}}}(1, 0, 0, 0, 0)^T$, (ii) $\Psi_U = \sqrt{n_{\text{tot}}}(0, 0, 1, 0, 0)^T$ or $\Psi_B = \sqrt{n_{\text{tot}}}(1, 0, 0, 0, 1)^T/\sqrt{2}$, and (iii) $\Psi_C = \sqrt{n_{\text{tot}}}(0, \sqrt{2}, 0, 0, 1)^T/\sqrt{3}$. In the mean-field theory, Ψ_B and Ψ_U are degenerate under zero magnetic field. It is predicted that zero point fluctuations lift this degeneracy [6]. The quadratic Zeeman effect also lifts this degeneracy, stabilizing Ψ_B for the $F = 2$ ^{87}Rb BEC at a magnetic field above the order of mG, which is consistent with experiments [8, 9, 10, 11]. We therefore do not consider the uniaxial nematic phase here.

The order parameter can be expressed in terms of the spherical harmonics: $\Phi(\theta, \phi) = \sum_{m=-2}^2 Y_{2m}(\theta, \phi) \psi_m$, where θ and ϕ are the polar and azimuthal angles in spin space, respectively. Figures 1 (a) and (b) show $\Phi(\theta, \phi)$ for the biaxial nematic phase Ψ_B and the cyclic phase Ψ_C . The profiles of $\Phi(\theta, \phi)$ for the biaxial nematic and cyclic phases are cloverleaf and triad, reflecting the D_4 and tetrahedral symmetries, respectively. Besides the trivial 2π gauge transformation and the 2π rotation in spin space, $\Phi(\theta, \phi)$ remains invariant under

some discrete spin-gauge transformations. In the biaxial nematic phase, for example, the combined operations of the π gauge transformation and the $\pi/2$ spin rotation around the z axis do not change $\Phi(\theta, \phi)$, namely $e^{i\pi} e^{-if^z \pi/2} \Psi_B = \Psi_B$. This discrete symmetry admits a fractional vortex around which the global gauge changes from 0 to π and the cloverleaf rotates by $\pi/2$. We refer to such a vortex as the 1/2–1/4 vortex, where 1/2 and 1/4 refer to the π gauge transformation and the $\pi/2$ spin rotation around the vortex, respectively. Similarly, we can define the 1/3–1/3 vortex in the cyclic phase [5, 7]. The order parameters around straight 1/2–1/4 and 1/3–1/3 vortices on the z axis can be expressed in cylindrical coordinates (r, φ, z) as

$$\Psi_{1/2-1/4}(r \rightarrow \infty) = e^{i\varphi/2} e^{-if^z \varphi/4} \Psi_B = \sqrt{\frac{n_{\text{tot}}}{2}} (1, 0, 0, 0, e^{i\varphi})^T, \quad (2)$$

$$\Psi_{1/3-1/3}(r \rightarrow \infty) = e^{i\varphi/3} e^{-if^z \varphi/3} \Psi_C = \sqrt{\frac{n_{\text{tot}}}{3}} (0, \sqrt{2}, 0, 0, e^{i\varphi})^T, \quad (3)$$

where the vortex is placed at $r = 0$. Among the vortices with nonzero mass circulation $\kappa = \hbar/M \oint d\mathbf{l} \sum_m \text{Im}[\psi_m^* (\nabla \psi_m)] / n_{\text{tot}}$, the 1/2–1/4 and 1/3–1/3 vortices are the most stable ones in the biaxial nematic and cyclic phases respectively [7].




FIG. 2: Schematic diagram of core structures for 1/2–1/4 (left) and 1/3–1/3 (right) vortices on the $\tilde{c}_1$ – $\tilde{c}_2$ plane. F: ferromagnetic core, B: biaxial nematic core, and C: cyclic core. The range of $(\tilde{c}_1, \tilde{c}_2)$ for a ^{87}Rb BEC is indicated [11].

Let us now examine the core structure. The core structure is determined by the competition among four terms in Eq. (1) and strongly depends on the interaction parameters $c_{0,1,2}$. We here consider the case for the 1/3–1/3 vortex in the cyclic phase: (i) When $c_{1,2} \gg c_0$, the third and fourth terms in Eq. (1) play a dominant role in determining the core structure. Because finite $\mathbf{F}$ or A_{20} drastically increases the energy, n_{tot} vanishes at the core. (ii) When $c_{1,2} \ll c_0$ and $c_1 < c_2$, on the contrary, n_{tot} is constant with the fixed total number of particles $N = \int d\mathbf{x} n_{\text{tot}}$ because of the inequality $\int d\mathbf{x} n_{\text{tot}}^2 \geq N^2$. A nonzero $\mathbf{F}$ raises energy less than A_{20} because of $c_1 < c_2$, and the core is ferromagnetic. (iii) When

$c_{1,2} \ll c_0$ and $c_1 > c_2$, the core is biaxial nematic for non-vanishing A_{20} with constant n_{tot} . (iv) When $c_{1,2} \sim c_0$, the last three parts in Eq. (1) compete. To minimize the kinetic energy, components with no phase singularity must be constant. Actually, $\psi_{m=2,1,0,-1}$ in Eq. (3) are constant inside and outside the core and the order parameter becomes $\sqrt{2n_{\text{tot}}/3}(0, 1, 0, 0, 0)$ at the ferromagnetic core. Similarly, we can show that the core of the $1/2-1/4$ vortex is cyclic for $|c_{1,2}| \ll c_0$ and $c_1 > 0$. Figure 2 shows a schematic diagram of the core structure in the $\tilde{c}_1 - \tilde{c}_2$ plane, where $\tilde{c}_1 \equiv c_1/c_0$ and $\tilde{c}_2 \equiv c_2/c_0$.

We next discuss the orientation of the order parameter of the $1/2-1/4$ vortex with the cyclic core that has to be embedded in the biaxial nematic order parameter. The energetically most favorable configuration of the order parameter at the core is found to be

$$\begin{aligned} \Psi_{1/2-1/4}(r=0) &= e^{2i\zeta/3} e^{-i\zeta f^z/3} e^{-if^x\pi} \Psi_C \\ &= \sqrt{n_{\text{tot}}/3}(1, 0, 0, \sqrt{2}e^{i\zeta}, 0)^T, \end{aligned} \quad (4)$$

where ζ is an arbitrary real number. Other possible candidates such as $e^{if^z\pi/4} e^{if^y \cos^{-1}(1/\sqrt{3})} \Psi_{1/2-1/4}(r=0) = (i/2, 0, 1/\sqrt{2}, 0, i/2)$ are energetically unfavorable, because it would require a drastic change from $\Psi_{1/2-1/4}(r \rightarrow \infty)$ to $\Psi_{1/2-1/4}(r=0)$. In Fig. 1 (c), we show $\Phi(\theta, \phi)$ for the $1/2-1/4$ vortex with the cyclic core. As shown in the figure, Eq. (4) fixes the order parameter of the core state in a manner such that the $(0, 0, 1)$ -directions in Figs. 1 (a) and (b) coincide with each other. As ζ changes, the order parameter shown in Fig. 1 (b) rotates around the z axis accompanied by a gauge transformation. Since the $m = -1$ component vanishes at $r \rightarrow \infty$, ζ does not contribute to the vortex energy. The remaining $SO(2)$ symmetry of the core state corresponding to ζ is spontaneously broken, leading to a localized Goldstone mode in the vortex core. For the $1/3-1/3$ vortex with the biaxial nematic core, we obtain the entire $\Phi(\theta, \phi)$ distribution as shown in Fig. 1 (d). In this case, the $(1, 1, 0)$ -direction in Fig. 1 (a) and the $(0, 0, 1)$ -direction in Fig. 1 (b) coincide with each other.

We now consider the ansatz for the $1/2-1/4$ vortex

$$\Psi_{1/2-1/4} = \sqrt{\frac{n_{\text{tot}}}{2}}(g, \alpha e^{i\varphi}, \beta e^{-i\varphi}, \gamma, f e^{i\varphi})^T, \quad (5)$$

where $\alpha, \beta, \gamma = 0$ and $f = g = 1$ at $r \rightarrow \infty$, and $\alpha = \beta = f = 0$, and $g = \sqrt{2/3}$ and $\gamma = 2/\sqrt{3}e^{i\zeta}$ at $r = 0$. Phase singularities for $m = 1, 0$ components are introduced to minimize the nonzero transverse magnetization [12]. The obtained transverse magnetization becomes: $F_x^2 + F_y^2 = n_{\text{tot}}|2(g\alpha^* + \gamma f^*) + \sqrt{6}(\beta\gamma^* + \alpha\beta^* e^{3i\varphi})|^2/6$, and the three-fold anisotropy arises. With this anisotropy, the Gross-Pitaevskii equation [4] derived from Eq. (1) also becomes anisotropic, leading to the anisotropic density distribution of each magnetic sublevel $|\psi_m|^2$. Similarly, the ansatz for the $1/3-1/3$ vortex which minimizes

the transverse magnetization is given by

$$\Psi_{1/3-1/3} = \sqrt{\frac{n_{\text{tot}}}{3}}(\alpha e^{i\varphi}, g, \beta e^{i\varphi}, \gamma, f e^{i\varphi})^T, \quad (6)$$

resulting in the two-fold anisotropy, where $\alpha, \beta, \gamma = 0$, $f = 1$, and $g = \sqrt{2}$ at $r \rightarrow \infty$, and $\alpha = \beta = f = 0$, $g = \sqrt{3/2}$, and $\gamma = \sqrt{3/2}e^{i\zeta}$ at $r = 0$. We have numerically minimized the mean-field energy (1) with the ansatz $\Psi_{1/2-1/4}$ and $\Psi_{1/3-1/3}$ with $\zeta = 0$. Figures 1 (e) and (f) show the obtained profiles of $|A_{30}|^2$ and $|A_{20}|^2$ for $\Psi_{1/2-1/4}$ and $\Psi_{1/3-1/3}$, respectively, where $A_{30} = -9(\psi_1^2\psi_{-2} + \psi_{-1}^2\psi_2) + \sqrt{6}\psi_0(-\psi_0^2 + 3\psi_1\psi_{-1} + 6\psi_2\psi_{-2})$ is the spin-singlet trio amplitude which is a measure of the cyclic order [2]. We can clearly see the triangular and elliptic deformations of the cores.

In Fig. 1 (c), the triad of the cyclic order parameter inside the core is embedded with the cloverleaf of the biaxial nematic order parameter, and there are three directions along which one of the three lobes of the triad is smoothly connected with one of two lobes of the cloverleaves surrounding the core. Along these directions, the triad is well overlapped with cloverleaves, which correspond to the minimum of $|\mathbf{F}|^2$ and the vertices of the triangle profile of $|A_{30}|^2$. Similarly, the biaxial nematic core of the $1/3-1/3$ vortex is deformed to be elliptic as shown in Fig. 1 (d). Thus, the triangular and elliptic shapes of the vortex core are the manifestations of the discrete three-fold and two-fold symmetries of the triad and the cloverleaf. Since the order parameter of the ferromagnetic phase does not have any discrete symmetry, vortices with the ferromagnetic core are isotropic.




FIG. 3: (Color) (a) $|A_{30}|^2$ and (b) $|\psi_0|^2$ of the $1/2-1/4$ vortex with the coupling constant $c_2^{(i)}$, and (c) $|A_{20}|^2$ and (d) $|\psi_2|^2$ of the $1/3-1/3$ vortex with $c_2^{(ii)}$. (e) $|A_{30}|^2$ of the $1/2-1/4$ vortex with $c_2^{(iii)}$. (f) $|A_{20}|^2$ of the $1/3-1/3$ vortex with $c_2^{(i)}$. Here a_{ho} is the harmonic oscillator length. $|A_{30}|^2$ and other values are normalized by $n_{\text{tot}}(r=0)^{3/2}$ and $n_{\text{tot}}(r=0)$, respectively.

We show that the vortices shown in Fig. 1 can be generated in the $F = 2$ ^{87}Rb BEC under an external rotation. For the $F = 2$ ^{87}Rb BEC, $\tilde{c}_1 \simeq (8.80 \pm 0.53) \times 10^{-3}$ and $\tilde{c}_2 \simeq (-4.71 \pm 5.15) \times 10^{-3}$ were experimentally ob-

tained [11], from which we can anticipate the 1/2–1/4 vortex with the cyclic core or the 1/3–1/3 vortex with the biaxial nematic core in this system (see Fig. 2). They have nonzero mass circulations $\kappa = h/2M$ and $h/3M$, respectively, and can be experimentally generated by an external rotation. In the presence of a trapping potential $V = M[\omega_r^2(x^2 + y^2) + \omega_z^2 z^2]/2$ and an external rotation $i\hbar\Omega_z(x\partial_y - y\partial_x)$ along the z axis, we find the stationary state of the order parameter for the $F = 2$ ^{87}Rb spinor BEC by minimizing the total energy. With experimental values of the radial frequency ω_r and the two-dimensional density [10], we perform two-dimensional simulation under $\Omega_z = 0.15\omega_r$. We use $c_{0,1,2}$ measured in [11]. For c_2 , we adopt the following two cases; $\tilde{c}_2^{(i)} = -4.71 \times 10^{-3}$ and $\tilde{c}_2^{(ii)} = 0.44 \times 10^{-3}$ where the ground state is expected to be biaxial nematic and cyclic, respectively. The stationary state can be obtained by calculating the imaginary-time development of the Gross-Pitaevskii equation derived from Eq. (1) [4]. We choose two initial states; stationary Ψ_B and Ψ_C without rotation for the 1/2–1/4 vortex and the 1/3–1/3 vortex, respectively. The results are shown in Figs. 3 (a) – (d). Both 1/2–1/4 and 1/3–1/3 vortices are stabilized under the rotation, and the spontaneous breaking of rotational symmetry also occurs reflecting the symmetry of the core state, which is shown in Fig. 3 (a) for $|A_{30}|^2$ in the 1/2–1/4 vortex and Fig. 3 (c) for $|A_{20}|^2$ in the 1/3–1/3 vortex. This symmetry breaking brings about an anisotropic density distribution, in particular, for the component described by α and β in Eqs. (5) and (6), and our results can be experimentally tested by the Stern-Gerlach experiment. In the present simulation, $|\psi_0|^2$ in the 1/2–1/4 vortex and $|\psi_2|^2$ in the 1/3–1/3 vortex are highly anisotropic as shown in Figs. 3 (b) and (d).

We have numerically confirmed that the vortex tiling phenomenon occurs independently of the sign of c_2 . Even when we start from the “false” ground state like Ψ_B with $c_2^{(ii)}$ or Ψ_C with $c_2^{(i)}$, those states involve the “metastable” 1/2–1/4 or 1/3–1/3 vortex, because the values of $c_{1,2}$ are so small that the spin relaxation dynamics from the false to true ground state is much slower than that of the vortex nucleation. Figures 3 (e) and (f) show the 1/2–1/4 vortex in Ψ_B with $c_2^{(ii)}$ and the 1/3–1/3 vortex in Ψ_C with $c_2^{(i)}$, and those figures are quite similar to Figs. 3 (a) and (c), respectively. These results imply that the nucleated vortex depends only on the initial state regardless of the sign of c_2 , and both cases of the 1/2–1/4 and 1/3–1/3 vortices can be experimentally realized using the $F = 2$ ^{87}Rb spinor BEC. In particular, the cyclic order parameter can be realized as the core of the 1/2–1/4 vortex surrounded by the biaxial nematic order parameter. Since both 1/2–1/4 and 1/3–1/3 vortices are non-Abelian [5, 7], the above method can be used to produce the first experimental realization of non-Abelian vortices.

The vortex tiling leads to a new type of excitation of vortices, that is a “twisting mode”. Due to the Goldstone mode with respect to ζ in Eq. (4), the triangular and elliptic vortex cores can rotate without energy cost. When we consider the 3-dimensional vortex line, there should be the characteristic wave excitation with a linear dispersion which corresponds to the twisting of the triangle or elliptic shape along the vortex line and can be defined as “twisting wave”. The detailed study of this mode will be reported elsewhere.

In conclusion, we have studied fractional vortices in a spin-2 spinor BEC and shown that the problem of vortex tiling in the order parameter space arises. A 1/2–1/4 vortex (1/3–1/3 vortex) with a cyclic core (biaxial nematic core) in the biaxial nematic (cyclic) phase requires the vortex tiling between the discrete D_4 symmetry of the biaxial nematic state and tetrahedral symmetry of the cyclic state. The rotational symmetry of the cores is spontaneously broken and their shapes become triangular for the 1/2–1/4 vortex and elliptic for the 1/3–1/3 vortex, which directly relate to the three-fold and two-fold rotational symmetries of the triad and cloverleaf structures. The vortex tiling can be experimentally implemented by using a rotating $F = 2$ ^{87}Rb BEC.

This work was supported by MEXT, Japan (KAKENHI, No. 207229 and 17071005, the Global COE Program “the Physical Sciences Frontier”, and the Photon Frontier Network Program).

-
- [1] J. Ruostekoski and J. R. Anglin, Phys. Rev. Lett. **91**, 190402 (2003).
 - [2] M. Koashi and M. Ueda, Phys. Rev. Lett. **84**, 1066 (2000); M. Ueda and M. Koashi, Phys. Rev. A **65**, 063602 (2002).
 - [3] C. V. Ciobanu, S.-K. Yip, and T.-L. Ho, Phys. Rev. A **61**, 033607 (2000).
 - [4] H. Saito and M. Ueda, Phys. Rev. A **72**, 053628 (2005).
 - [5] S. -K. Yip, Phys. Rev. A **75**, 023625 (2007).
 - [6] J. L. Song, G. W. Semenoff, and F. Zhou, Phys. Rev. Lett. **98**, 160408 (2007); A. M. Turner, *et al.*, Phys. Rev. Lett. **98**, 190404 (2007).
 - [7] H. Mäkelä, Y. Zhang and K.-A. Suominen, J. Phys. A: Math. Gen. **36**, 8555 (2003); H. Mäkelä, J. Phys. A: Math. Gen. **39**, 7423 (2006); G. W. Semenoff and F. Zhou, Phys. Rev. Lett. **98**, 100401 (2007); M. Kobayashi *et al.*, arXiv:0810.5441 (2008).
 - [8] H. Schmaljohann *et al.*, Phys. Rev. Lett. **92**, 040402 (2004).
 - [9] M.-S. Chang *et al.*, Phys. Rev. Lett. **92**, 140403 (2004).
 - [10] T. Kuwamoto *et al.*, Phys. Rev. A **69**, 063604 (2004); Tojo *et al.*, Appl. Phys. B **93**, 403 (2008).
 - [11] A. Widera *et al.*, New J. Phys. **8**, 152 (2006).
 - [12] M. Kobayashi, Y. Kawaguchi, and M. Ueda (to be published).